Chief, I a S Staff

A/B, II, 9, 47

Bluebird Team

Modical Summary File No.: B/2 - 1, 2, 3, 4

The drug used in this case was Sedius Acytal intravenessly, 5% solution given at the rate of 1 or a minute. The total wount absintance was 16 or or 0.8 ms. The shi when was Dencedrine, 20 mg, 2-5 or intravenessly, A total of 9 or was given or 36 mg.

The injection was started at 1141. At 1152, after the administration of 11 cc of Sodium Amytal solution, the Subject said his head folt weary. At 1155, he was morely acloud but arrichaged with a start upon being arounds. From that time on, he was scansionally given 1 cc more of the solution in order to anintain the proper state for the technique. At 1250, he was again esleep but easily around and the same at 1245.

The desired effect of the medication was obtained. The Subject was violent upon occanions during the transference, at which time, he was convinced that the interrogators were the subject was begun and although he gradually awakened, at 1259, after the administration of 7 cc of Bonzedrine solution, he sail that he still felt sleepy. He was given 2 cc more at 1303. At 1553, he was awake. At 1404, he was sent to eat and then sleep. He said he still felt a limble diszy.

The desired effect of the medication proviously mentioned was obtained. The Subject had no suspicion of other than the use of the medication to help him in connection with the 0. F. machine.

STONED

___ - A

1 (
12 miles of the second
8-11
1
H-4, 15
7. 12
. s.
11. 15715
START UECK LEVY
REAKLY LELETT, EUT
17
D.D. Marke - RESERVETA File Sala CENTER
1 1
1
1、11

-

1354